

## Random walks on graphs

Most of the material in this handout is taken from: Daniel Spielman, *Spectral and Algebraic Graph Theory. Manuscript, 2019.*

Let  $A$  be the adjacency matrix of a graph  $G$ . We use  $\alpha_1 \geq \dots \geq \alpha_n$  to denote the eigenvalues of  $A$  (note that they are ordered in the opposite direction with respect to the eigenvalues  $\lambda_1 \leq \dots \leq \lambda_n$  of the Laplacian matrix  $L$ ).

If  $G$  is  $d$ -regular, then  $L = I - \frac{1}{d}A$  and therefore  $\lambda_i = 1 - \frac{\alpha_i}{d}$ . Since  $\lambda_i \in [0, 2]$  for any  $G$  (even not regular), we have that  $\alpha_i \in [-d, d]$  for any  $d$ -regular graph.

**Fact 1** For any graph  $G = (V, E)$ ,  $d(G) \leq \alpha_1 \leq \Delta(G)$ .

PROOF. Using the variational characterization of eigenvalues,

$$\alpha_1 = \max_{\mathbf{x}: \mathbf{x} \neq \mathbf{0}} \frac{\mathbf{x}^\top A \mathbf{x}}{\mathbf{x}^\top \mathbf{x}} \geq \frac{\mathbf{1}^\top A \mathbf{1}}{\mathbf{1}^\top \mathbf{1}} = \frac{1}{n} \sum_{i=1}^n \sum_{j=1}^n A_{i,j} = \frac{1}{n} \sum_{i=1}^n d(i) = d(G)$$

For the other inequality, let  $\mathbf{u}$  be an eigenvector for the eigenvalue  $\alpha_1$  and let  $u_i > 0$  the largest component of  $\mathbf{u}$  (if all components are negative, take  $-\mathbf{u}$ ). Then

$$\alpha_1 = \frac{(A\mathbf{u})_i}{u_i} = \frac{1}{u_i} \sum_{j=1}^n A_{i,j} u_j = \sum_{j=1}^n A_{i,j} \frac{u_j}{u_i} \leq \sum_{j=1}^n A_{i,j} \leq \Delta(G)$$

concluding the proof. □

The trace of a symmetric  $n \times n$  matrix  $M$  is  $M_{1,1} + \dots + M_{n,n}$ . One can show that the trace is equal to the sum of eigenvalues. Since  $A_{i,i} = 0$ , the trace of  $A$  is zero and so  $\alpha_1 + \dots + \alpha_n = 0$ . Since we proved that  $\alpha_1 > d(G) > 0$ , this implies that  $\alpha_n < 0$ .

**Lemma 2** Let  $G = (V, E)$  be a connected graph and let  $M$  be a nonnegative symmetric matrix such that  $M_{i,j} > 0$  if and only if  $(i, j) \in E$ . Assume that some nonnegative vector  $\mathbf{u}$  is an eigenvector of  $M$ . Then  $\mathbf{u}$  is strictly positive.

PROOF. If  $\mathbf{u}$  is nonnegative but not strictly positive, then there is some vertex  $r$  for which  $u_r = 0$ . As  $G$  is connected, there must be some edge  $(r, s)$  for which  $u_r = 0$  but  $u_s > 0$  (since  $\mathbf{u}$  is an eigenvector,  $\mathbf{u} \neq \mathbf{0}$ ). Let  $\mu$  be the eigenvalue of  $\mathbf{u}$ . We obtain a contradiction from

$$0 = \mu u_r = (M\mathbf{u})_r = \sum_{i=1}^n M_{r,i} u_i \geq M_{r,s} u_s > 0$$

concluding the proof. □

The next result is the cornerstone for the analysis of random walks on graphs. It applies to many symmetric matrices defined on graphs, including the adjacency matrix and the normalized adjacency matrix..

**Theorem 3 (Perron-Frobenius for symmetric matrices)** Let  $G = (V, E)$  be a connected graph and let  $M$  be a nonnegative symmetric matrix such that, for all  $i \neq j$ ,  $M_{i,j} > 0$  if and only if  $(i, j) \in E$ . Then the eigenvalues  $\mu_1 \geq \dots \geq \mu_n$  of  $M$  satisfy:

1. The largest eigenvalue  $\mu_1$  has a strictly positive eigenvector,
2.  $\mu_1 \geq -\mu_n$ ,
3.  $\mu_1 > \mu_2$ , implying that  $\mu_1$  has multiplicity 1.

PROOF. In order to prove part 1, let  $\mathbf{u}_1$  an eigenvector for  $\mu_1$  and define  $x_i = |u_{1,i}|$ . Then  $\mathbf{x}$  is also an eigenvector. Indeed,  $\mathbf{x}^\top \mathbf{x} = \mathbf{u}_1^\top \mathbf{u}_1 = 1$ . Moreover, since  $M$  is nonnegative,

$$\mu_1 = \mathbf{u}_1^\top M \mathbf{u}_1 = \sum_{i=1}^n \sum_{j=1}^n M_{i,j} u_i u_j \leq \sum_{i=1}^n \sum_{j=1}^n M_{i,j} |u_i| |u_j| = \mathbf{x}^\top M \mathbf{x}$$

Since  $\mu_1$  is the largest eigenvalue of  $M$ ,  $\mathbf{x}$  must be an eigenvector of  $\mu_1$ . Since  $\mathbf{x}$  is nonnegative, Lemma 2 implies that is strictly positive.

To prove part 2, let  $\mathbf{u}_n$  be an eigenvector of  $\mu_n$  and let  $x_i = |u_{n,i}|$ . Then, similarly to before and recalling that  $\mu_n < 0$ ,

$$|\mu_n| = |\mathbf{u}_n^\top M \mathbf{u}_n| \leq \sum_{i=1}^n \sum_{j=1}^n M_{i,j} |u_i| |u_j| = \mathbf{x}^\top M \mathbf{x} \leq \mu_1$$

To prove part 3, consider an eigenvector  $\mathbf{u}_2$  of  $\mu_2$ . Since  $\mathbf{u}_2^\top \mathbf{u}_1 = 0$ ,  $\mathbf{u}_2$  must contain positive and negative components. Now let  $x_i = |u_{2,i}|$  and, once again, note that  $\mu_2 = \mathbf{u}_2^\top M \mathbf{u}_2 \leq \mathbf{x}^\top M \mathbf{x} \leq \mu_1$ . Since  $\mathbf{u}_2$  has positive and negative components and the graph is connected, there must be at least one edge  $(i, j) \in E$  such that  $u_{2,i} < 0 < u_{2,j}$ . This edge gives a negative contribution to  $\mathbf{u}_2^\top M \mathbf{u}_2$  and a positive contribution to  $\mathbf{x}^\top M \mathbf{x} \leq \mu_1$  (recall that  $M$  is nonnegative). Hence the inequality  $\mathbf{u}_2^\top M \mathbf{u}_2 \leq \mathbf{x}^\top M \mathbf{x}$  must be strict, implying  $\mu_2 < \mu_1$ .  $\square$

The next observation (proof omitted) is important in the analysis of convergence of a random walk on a graph.

**Fact 4**  $G$  is bipartite if and only if  $\mu_n = -\mu_1$ .

**The random walk on a graph.** Given a connected graph  $G = (V, E)$  with  $V = \{1, \dots, n\}$ , we consider the random walk that starts from an arbitrary vertex  $V_0 \in V$ , and at each step  $t = 0, 1, \dots$  moves from  $V_t$  to a random vertex  $V_{t+1}$  in the neighborhood of  $V_t$ . Let  $\mathbf{e}_i$  be the canonical basis vector for the  $i$ -th coordinate (all zeros but a single 1 in position  $i$ ). The state of the walk at time  $t$  is defined by a probability distribution  $\mathbf{p}_t$  over  $V$ . Hence, if the walk starts at  $V_0 = i$ , then  $\mathbf{p}_0 = \mathbf{e}_i$ . At any time  $t$  we have

$$\mathbf{p}_t(i) = \mathbb{P}(V_t = i) = \sum_{j: (i,j) \in E} \mathbb{P}(V_t = i \mid V_{t-1} = j) \mathbb{P}(V_{t-1} = j) = \sum_{j=1}^n \frac{A_{i,j}}{d(j)} \mathbf{p}_{t-1}(j) \quad (1)$$

Let  $D = \text{diag}(d(1), \dots, d(n))$  and note that  $D_{i,j}^{-1} = \mathbb{I}\{i = j\}/d(j)$ . Since

$$(AD^{-1})_{i,j} = \sum_{k=1}^n A_{i,k} \frac{\mathbb{I}\{k = j\}}{d(j)} = \frac{A_{i,j}}{d(j)}$$

the right-hand side of (1) can be rewritten as  $AD^{-1}\mathbf{p}_{t-1}$ . Letting  $W = AD^{-1}$ , the evolution of our random walk is given by  $\mathbf{p}_t = W\mathbf{p}_{t-1}$ , or  $\mathbf{p}_t = W^t\mathbf{p}_0$ .  $W$  is a column-stochastic matrix, as it is a nonnegative matrix whose elements in each column sum to 1, that is  $\mathbf{1}^\top W = \mathbf{1}^\top$ .

As  $W_{i,j} = A_{i,j}/d(j)$ , the matrix  $W$  is not symmetric. However, it is related to the normalized adjacency matrix  $A_{\text{norm}} = D^{-1/2}AD^{-1/2}$  (which is symmetric), indeed,  $A_{\text{norm}} = D^{-1/2}WD^{1/2}$ . The normalized adjacency matrix is in turn related to the normalized Laplacian as follows

$$L_{\text{norm}} = I - D^{-1/2}AD^{-1/2} = I - A_{\text{norm}}$$

Let  $\omega_1 \geq \dots \geq \omega_n$  be the eigenvalues of  $A_{\text{norm}}$ . Because the eigenvalues  $\lambda_1 \leq \dots \leq \lambda_n$  of  $L_{\text{norm}}$  belong to the interval  $[0, 2]$ , it is easy to verify that  $\omega_i = 1 - \lambda_i$ . This immediately implies that  $\omega_i \in [-1, 1]$  for all  $i$ .

**Fact 5** *The vector  $\boldsymbol{\psi}$  is an eigenvector of  $A_{\text{norm}}$  of eigenvalue  $\omega$  if and only if  $D^{1/2}\boldsymbol{\psi}$  is an eigenvector of  $W$  of eigenvalue  $\omega$ .*

PROOF. As  $A_{\text{norm}} = D^{-1/2}WD^{1/2}$ , we have that  $D^{1/2}A_{\text{norm}} = WD^{1/2}$ . Thus, if  $A_{\text{norm}}\boldsymbol{\psi} = \omega\boldsymbol{\psi}$ , then

$$WD^{1/2}\boldsymbol{\psi} = D^{1/2}A_{\text{norm}}\boldsymbol{\psi} = D^{1/2}\omega\boldsymbol{\psi} = \omega(D^{1/2}\boldsymbol{\psi})$$

and, similarly, we can show that  $W\mathbf{u} = \omega\mathbf{u}$  implies  $A_{\text{norm}}(D^{-1/2}\mathbf{u}) = \omega(D^{-1/2}\mathbf{u})$ .  $\square$

This result implies that  $W$  has eigenvalues  $\omega_i \in [-1, 1]$ .

An application of the Perron-Frobenius theorem to the normalized adjacency matrix  $A_{\text{norm}}$  gives:

1. the largest eigenvalue  $\omega_1$  has a unique eigenvector  $\boldsymbol{\psi}_1$  whose components are strictly positive
2.  $\omega_n = -\omega_1$  if and only if  $G$  is bipartite.

**The stationary distribution.** We say that a distribution  $\boldsymbol{\pi}$  over  $V$  is the stationary distribution of  $W$  if  $W\boldsymbol{\pi} = \boldsymbol{\pi}$ . Hence, the stationary distribution is a (unnormalized) eigenvector of  $W$  with eigenvalue  $1 = \omega_1$ , as the eigenvalues of  $W$  range in  $[-1, 1]$ . Now let  $\mathbf{d} = (d(1), \dots, d(n))$  be the vector of vertex degrees and consider the distribution

$$\mathbf{p} = \frac{\mathbf{d}}{\mathbf{1}^\top \mathbf{d}}$$

Since

$$(W\mathbf{p})_i = \frac{1}{\mathbf{1}^\top \mathbf{d}} \sum_{j=1}^n A_{i,j} \left( \sum_{k=1}^n D_{j,k}^{-1} d(k) \right) = \frac{1}{\mathbf{1}^\top \mathbf{d}} \sum_{j=1}^n A_{i,j} = \frac{d(i)}{\mathbf{1}^\top \mathbf{d}} = \mathbf{p}(i)$$

we have that  $\mathbf{p}$  is the stationary distribution  $\boldsymbol{\pi}$  for  $W$ . Moreover, we also know that  $\boldsymbol{\pi}$  is (up to normalization) the unique eigenvector of  $W$  for the eigenvalue 1. So Fact 5 implies  $\boldsymbol{\pi} = D^{1/2}\boldsymbol{\psi}_1$ .

**Fact 6** *Let  $A$  be a  $n \times n$  symmetric matrix with spectrum  $\lambda_1, \dots, \lambda_n, \mathbf{u}_1, \dots, \mathbf{u}_n$ . Then, for any  $t \in \mathbb{N}$ ,*

$$A^t = \sum_{i=1}^n \lambda_i^t \mathbf{u}_i \mathbf{u}_i^\top$$

PROOF. We use induction on  $t$  together with the spectral theorem and the orthonormality of the eigenvectors. For  $t = 1$  the statement follows from the spectral theorem. Assume now the statement holds for  $t - 1$  and write

$$\begin{aligned}
A^t &= AA^{t-1} = \left( \sum_{i=1}^n \lambda_i \mathbf{u}_i \mathbf{u}_i^\top \right) \left( \sum_{j=1}^n \lambda_j^{t-1} \mathbf{u}_j \mathbf{u}_j^\top \right) \\
&= \sum_{i=1}^n \sum_{j=1}^n \lambda_i \lambda_j^{t-1} \mathbf{u}_i \mathbf{u}_i^\top \mathbf{u}_j \mathbf{u}_j^\top \\
&= \sum_{i=1}^n \lambda_i^t \mathbf{u}_i \mathbf{u}_i^\top \quad (\text{since } \mathbf{u}_i^\top \mathbf{u}_j = \mathbb{I}\{i = j\})
\end{aligned}$$

concluding the proof.  $\square$

We are now ready to prove the convergence of the random walk to the stationary distribution.

**Theorem 7** *For any connected graph  $G$  not bipartite,*

$$\lim_{t \rightarrow \infty} W^t \mathbf{p}_0 = \frac{\mathbf{d}}{\mathbf{1}^\top \mathbf{d}}$$

*irrespective to the initial distribution  $\mathbf{p}_0$ .*

PROOF. To verify convergence to  $\boldsymbol{\pi}$ , we express  $D^{-1/2} \mathbf{p}_0$  in the eigenbasis  $\boldsymbol{\psi}_1, \dots, \boldsymbol{\psi}_n$  of  $A_{\text{norm}}$ ,

$$D^{-1/2} \mathbf{p}_0 = \sum_{i=1}^n \left( \boldsymbol{\psi}_i^\top D^{-1/2} \mathbf{p}_0 \right) \boldsymbol{\psi}_i = \sum_{i=1}^n c_i \boldsymbol{\psi}_i \quad (2)$$

Now we write

$$\begin{aligned}
\mathbf{p}_t &= W^t \mathbf{p}_0 = \left( D^{1/2} A_{\text{norm}} D^{-1/2} \right)^t \mathbf{p}_0 \\
&= D^{1/2} A_{\text{norm}} D^{-1/2} D^{1/2} A_{\text{norm}} D^{-1/2} \dots D^{1/2} A_{\text{norm}} D^{-1/2} \mathbf{p}_0 \\
&= D^{1/2} A_{\text{norm}}^t D^{-1/2} \mathbf{p}_0 \quad (\text{using } D^{-1/2} D^{1/2} = I) \\
&= D^{1/2} A_{\text{norm}}^t \sum_{i=1}^n c_i \boldsymbol{\psi}_i \quad (\text{using (2)}) \\
&= D^{1/2} \left( \sum_{j=1}^n \omega_j^t \boldsymbol{\psi}_j \boldsymbol{\psi}_j^\top \right) \sum_{i=1}^n c_i \boldsymbol{\psi}_i \quad (\text{using Fact 6}) \\
&= D^{1/2} \sum_{i=1}^n \sum_{j=1}^n c_i \omega_j^t \boldsymbol{\psi}_j \boldsymbol{\psi}_j^\top \boldsymbol{\psi}_i \\
&= D^{1/2} \sum_{i=1}^n c_i \omega_i^t \boldsymbol{\psi}_i \quad (\text{since } \boldsymbol{\psi}_i^\top \boldsymbol{\psi}_j = \mathbb{I}\{i = j\})
\end{aligned}$$

Therefore, recalling that  $\omega_1 = 1$ ,

$$\mathbf{p}_t = D^{1/2}c_1\boldsymbol{\psi}_1 + D^{1/2}\sum_{i=2}^n c_i\omega_i^t\boldsymbol{\psi}_i \quad (3)$$

Now, if  $G$  is not bipartite, then  $\omega_2, \dots, \omega_n \in (-1, 1)$ . Since  $\lim_{x \rightarrow \infty} \omega^x = 0$  for all  $\omega \in (-1, 1)$ , we get

$$\lim_{t \rightarrow \infty} \mathbf{p}_t = D^{1/2}c_1\boldsymbol{\psi}_1$$

Now recall that  $\boldsymbol{\pi}$  is a unnormalized eigenvector of  $W$ . Therefore, using Fact 5,  $\boldsymbol{\pi} = \frac{\mathbf{d}}{\mathbf{1}^\top \mathbf{d}} \propto D^{1/2}\boldsymbol{\psi}_1$  implying  $\boldsymbol{\psi}_1 = D^{-1/2}\mathbf{d}/\|D^{-1/2}\mathbf{d}\|$ . Therefore

$$c_1 = \boldsymbol{\psi}_1^\top D^{-1/2}\mathbf{p}_0 = \frac{(D^{-1/2}\mathbf{d})^\top}{\|D^{-1/2}\mathbf{d}\|} D^{-1/2}\mathbf{p}_0 = \frac{\mathbf{d}^\top D^{-1}\mathbf{p}_0}{\|D^{-1/2}\mathbf{d}\|} = \frac{\mathbf{1}^\top \mathbf{p}_0}{\|D^{-1/2}\mathbf{d}\|} = \frac{1}{\|D^{-1/2}\mathbf{d}\|}$$

because  $\mathbf{p}_0$  is a probability vector. So,

$$D^{1/2}c_1\boldsymbol{\psi}_1 = \frac{1}{\|D^{-1/2}\mathbf{d}\|} D^{1/2}D^{-1/2} \frac{\mathbf{d}}{\|D^{-1/2}\mathbf{d}\|} = \frac{\mathbf{d}}{\|D^{-1/2}\mathbf{d}\|^2} = \frac{\mathbf{d}}{\sum_{j=1}^n d(j)^2/d(j)} = \boldsymbol{\pi}$$

concluding the proof.  $\square$

**Speed of convergence of the random walk.** Assume that the random walk starts at some vertex  $u \in V$ . For every vertex  $v \in V$ , we will bound how far  $\mathbf{p}_t(v)$  can be from  $\boldsymbol{\pi}(v)$ .

**Theorem 8** *For all  $u, v \in V$  and  $t \in \mathbb{N}$ , if  $\mathbf{p}_0 = \mathbf{e}_u$ , then*

$$|\mathbf{p}_t(v) - \boldsymbol{\pi}(v)| \leq \left( \sqrt{\frac{d(v)}{d(u)}} \right) \kappa^t$$

where  $\kappa = \left( \max \{ |\omega_n|, |\omega_2| \} \right)$ .

PROOF. We start by writing  $\mathbf{p}_t(v) = \mathbf{e}_v^\top \mathbf{p}_t$ . Recalling (3),

$$\mathbf{p}_t(v) = \mathbf{e}_v^\top \mathbf{p}_t = \boldsymbol{\pi}(v) + \mathbf{e}_v^\top D^{1/2} \sum_{i=2}^n \omega_i^t c_i \boldsymbol{\psi}_i \quad (4)$$

Using (2), we know that

$$c_i = \boldsymbol{\psi}_i^\top D^{-1/2} \mathbf{e}_u = \frac{\boldsymbol{\psi}_i^\top \mathbf{e}_u}{\sqrt{d(u)}}$$

So, from (4) and  $\mathbf{e}_v^\top D^{1/2} = \sqrt{d(v)} \mathbf{e}_v^\top$ ,

$$\mathbf{e}_v^\top D^{1/2} \sum_{i=2}^n \omega_i^t c_i \boldsymbol{\psi}_i = \left( \sqrt{\frac{d(v)}{d(u)}} \right) \mathbf{e}_v^\top \sum_{i=2}^n \omega_i^t \boldsymbol{\psi}_i \boldsymbol{\psi}_i^\top \mathbf{e}_u$$

Now we look at the last part of the above expression. We can write

$$\begin{aligned}
\mathbf{e}_v^\top \sum_{i=2}^n \omega_i^t \boldsymbol{\psi}_i \boldsymbol{\psi}_i^\top \mathbf{e}_u &= \sum_{i=2}^n \omega_i^t (\mathbf{e}_v^\top \boldsymbol{\psi}_i) (\boldsymbol{\psi}_i^\top \mathbf{e}_u) \\
&\leq \sum_{i=2}^n |\omega_i|^t |\mathbf{e}_v^\top \boldsymbol{\psi}_i| |\boldsymbol{\psi}_i^\top \mathbf{e}_u| \\
&\leq \kappa^t \sum_{i=1}^n |\mathbf{e}_v^\top \boldsymbol{\psi}_i| |\boldsymbol{\psi}_i^\top \mathbf{e}_u| \\
&\leq \kappa^t \sqrt{\sum_{i=1}^n (\mathbf{e}_v^\top \boldsymbol{\psi}_i)^2} \sqrt{\sum_{i=1}^n (\boldsymbol{\psi}_i^\top \mathbf{e}_u)^2} \quad (\text{using the Cauchy-Schwartz inequality}) \\
&= \kappa^t \|\mathbf{e}_v\| \|\mathbf{e}_u\| \quad (\text{because } \boldsymbol{\psi}_1, \dots, \boldsymbol{\psi}_n \text{ is an orthonormal basis}) \\
&= \kappa^t
\end{aligned}$$

This concludes the proof.  $\square$

**The lazy random walk.** Note that  $\omega_2$  can be negative. For instance, if  $G = K_n$ , then  $\omega_2 = \dots = \omega_n = -\frac{1}{n-1}$ . In order to bring out the main message in the analysis of the convergence time, we can replace  $W$  by

$$W' = \frac{1}{2}(I + W)$$

With this new matrix, with equal probabilities we have that  $V_{t+1} = V_t$  or  $V_{t+1}$  is a random neighbor of  $V_t$ . Also the eigenvalues  $\omega'_1 \geq \dots \geq \omega'_n$  of  $W'$  satisfy  $\omega'_i = \frac{1}{2}(1 + \omega_i) \in [0, 1]$ . It is easy to check that  $\boldsymbol{\pi} = \frac{\mathbf{d}}{\mathbf{1}^\top \mathbf{d}}$  is the stable distribution also for  $W'$ . The relation between  $W'$  and  $L_{\text{norm}}$  is now

$$W' = I - \frac{1}{2} D^{1/2} L_{\text{norm}} D^{-1/2}$$

Doing again the proof of Theorem 8 we obtain

$$\mathbf{p}_t(v) = \boldsymbol{\pi} + D^{1/2} \sum_{i=2}^n c_i \left(1 - \frac{\lambda_i}{2}\right)^t \boldsymbol{\psi}_i$$

where  $1 - \frac{\lambda_i}{2} = \omega'_i$ . Therefore, when applying Theorem 8 to  $W'$  we find that  $\kappa = \omega'_2$ .

**Mixing time.** The mixing time of  $W'$  is the smallest  $t$  such that

$$|(W')^t \mathbf{e}_u - \boldsymbol{\pi}(v)| \leq \frac{\boldsymbol{\pi}(v)}{2} \quad \text{for all } u, v \in V$$

Since  $\kappa = \omega_2 = 1 - \frac{\lambda_2}{2}$ , where  $\lambda_2$  is the smallest nonzero eigenvalue of the normalized Laplacian,  $W$  has mixed when

$$\left(\sqrt{\frac{d(v)}{d(u)}}\right) \left(1 - \frac{\lambda_2}{2}\right)^t \leq \frac{d(v)}{2\mathbf{1}^\top \mathbf{d}} \quad \text{for all } v \in V$$

Using  $2|E| = \mathbf{1}^\top \mathbf{d}$  and  $1 + x \leq e^x$  for all  $x \in \mathbb{R}$  the above is equivalent to

$$t \geq \frac{2}{\lambda_2} \ln \frac{4|E|}{\sqrt{d(u)d(v)}} \quad \text{which is implied by} \quad t \geq \frac{2}{\lambda_2} \ln \frac{4|E|}{\delta(G)}$$

This reveals the important connection between mixing time and clusterability and the role of  $\lambda_2$ .

**Distributed consensus.** Given numbers  $\mathbf{x}_0 = (\mathbf{x}_0(1), \dots, \mathbf{x}_0(n))$  at each vertex of a connected graph  $G = (V, E)$ , we want each node in  $V$  to compute the average

$$\mu = \frac{1}{n} \sum_{v \in V} \mathbf{x}_0(v)$$

by communicating only with its neighbors in  $G$ .

We run at each node  $v \in V$  an algorithm that, at each time step  $t = 0, 1, \dots$ , updates the node's state  $\mathbf{x}_t(v)$  according to

$$\mathbf{x}_{t+1}(v) = \sum_{u: (u,v) \in E} W_{u,v} \mathbf{x}_t(u) \quad (5)$$

We can write the update as  $\mathbf{x}_{t+1} = W\mathbf{x}_t$ . We let the matrix  $W$  to be a **gossip matrix**. This is any nonnegative symmetric matrix, doubly stochastic ( $W\mathbf{1} = \mathbf{1}$  and  $\mathbf{1}^\top W = \mathbf{1}^\top$ ), and such that  $W_{i,j} > 0$  if and only if  $(i, j) \in E$ .

**Fact 9** *The largest eigenvalue of a row-stochastic matrix is 1.*

PROOF. Let  $W$  be a row-stochastic matrix. Then  $W\mathbf{1} = \mathbf{1}$  and so 1 is an eigenvalue of  $W$ . Now suppose there exists  $\mu > 1$  and  $\mathbf{x} \neq \mathbf{0}$  such that  $W\mathbf{x} = \mu\mathbf{x}$ . Let  $x_k$  be a largest element of  $\mathbf{x}$ . Since any scalar multiple of  $\mathbf{x}$  will also satisfy this equation we can assume, without loss of generality, that  $x_k > 0$ . Since the elements  $W_{i,1}, \dots, W_{i,n}$  on each row of  $W$  are nonnegative and sum to 1, for any  $i = 1 \dots, n$  we have

$$(W\mathbf{x})_i = \sum_{j=1}^n W_{i,j} x_j \leq \max_{j=1, \dots, n} x_j = x_k$$

Thus, no entry in  $\mu\mathbf{x} = W\mathbf{x}$  can be larger than  $x_k$ . But since  $\mu > 1$ ,  $\mu x_k > x_k$  and we have a contradiction. Therefore, the largest eigenvalue of  $W$  is 1.  $\square$

Let  $\omega_n \leq \dots \leq \omega_{n-1} < \omega_1 = 1$  be the eigenvalues of  $W$ . Let

$$\boldsymbol{\mu} = \mu\mathbf{1} = \frac{1}{n} \mathbf{1}\mathbf{1}^\top \mathbf{x}_0$$

Then  $\boldsymbol{\mu}$  is the stationary distribution for  $W$ . Indeed, because  $W$  is row-stochastic,

$$W\boldsymbol{\mu} = \frac{1}{n} W\mathbf{1}\mathbf{1}^\top \mathbf{x}_0 = \frac{1}{n} \mathbf{1}\mathbf{1}^\top \mathbf{x}_0 = \boldsymbol{\mu}$$

Note also that

$$\frac{1}{n} \mathbf{1}\mathbf{1}^\top \mathbf{x}_t = \frac{1}{n} \mathbf{1}\mathbf{1}^\top W\mathbf{x}_{t-1} = \frac{1}{n} \mathbf{1}\mathbf{1}^\top \mathbf{x}_{t-1} = \dots = \frac{1}{n} \mathbf{1}\mathbf{1}^\top \mathbf{x}_0 = \frac{1}{n} \mathbf{1}\mathbf{1}^\top \boldsymbol{\mu} \quad (6)$$

In order to prove convergence of  $\mathbf{x}_t$  to  $\boldsymbol{\mu}$ , we measure how fast  $\|\mathbf{x}_t - \boldsymbol{\mu}\|$  vanishes as  $t \rightarrow \infty$ . We use the operator norm  $\|W\|$  of a symmetric matrix  $W$ , which is the absolute value of the largest eigenvalue of  $W$ . For any vector  $\mathbf{x}$ , we have the following inequality:  $\|W\mathbf{x}\| \leq \|W\| \|\mathbf{x}\|$ . Using that inequality and (6) we can write

$$\|\mathbf{x}_{t+1} - \boldsymbol{\mu}\|^2 = \|W(\mathbf{x}_t - \boldsymbol{\mu})\|^2 = \left\| \left( W - \frac{\mathbf{1}\mathbf{1}^\top}{n} \right) (\mathbf{x}_t - \boldsymbol{\mu}) \right\|^2 \leq \left\| W - \frac{\mathbf{1}\mathbf{1}^\top}{n} \right\|^2 \|\mathbf{x}_t - \boldsymbol{\mu}\|^2$$

**Fact 10** *If  $G$  is not bipartite, then*

$$\left\| W - \frac{\mathbf{1}\mathbf{1}^\top}{n} \right\| = |\omega_2|$$

PROOF. By Fact 9,  $\omega_1 = 1$  with eigenvalue  $\mathbf{u}_1 = \frac{1}{\sqrt{n}}\mathbf{1}$ . Let  $W = U\Lambda U^\top$  be the spectral decomposition of  $W$ , where  $\Lambda = \text{diag}(1, \omega_2, \dots, \omega_n)$  and  $U = [\mathbf{u}_1, \dots, \mathbf{u}_n]$ . Let  $M$  be the  $n \times n$  diagonal matrix  $\text{diag}(1, 0, \dots, 0)$ . Then,  $UMU^\top = \mathbf{u}_1\mathbf{u}_1^\top = \frac{1}{n}\mathbf{1}\mathbf{1}^\top$  and

$$\left\| W - \frac{\mathbf{1}\mathbf{1}^\top}{n} \right\| = \left\| U(\Lambda - M)U^\top \right\| = \left\| U \text{diag}(0, \omega_2, \dots, \omega_n)U^\top \right\| = |\omega_2|$$

because  $G$  is not bipartite and so  $\omega_2$  is the largest eigenvalue of the matrix  $U \text{diag}(0, \omega_2, \dots, \omega_n)U^\top$ .  
□

Hence  $\|\mathbf{x}_t - \boldsymbol{\mu}\| \leq |\omega_2|^t \|\mathbf{x}_0 - \boldsymbol{\mu}\|$ , implying that the speed of convergence is dictated by the second largest eigenvalue of the gossip matrix  $W$  if the underlying graph is not bipartite.

A reasonable choice for the gossip matrix is  $W = I - \alpha L$ , where  $\alpha > 0$  and  $L = D - A$  is the unnormalized Laplacian of  $G$  (check that this choice of  $W$  is indeed a gossip matrix). The eigenvalues of  $W$  are thus  $\omega_i = 1 - \alpha\lambda_i$ , where  $\alpha \leq 1/\Delta(G)$  to respect the nonnegativity condition for the entries of  $W$  and  $0 = \lambda_1 < \lambda_2 \leq \dots \leq \lambda_n$  are the eigenvalues of  $L$ . The update (5) in this case can be written as

$$\mathbf{x}_{t+1}(v) = \mathbf{x}_t(v) + \alpha \sum_{u:(u,v) \in E} (\mathbf{x}_t(u) - \mathbf{x}_t(v))$$

and the speed of convergence is dictated by  $\lambda_2/\Delta(G)$ ,

$$|\omega_2|^t \leq \left(1 - \frac{\lambda_2}{\Delta(G)}\right)^t \leq \exp\left(-t \frac{\lambda_2}{\Delta(G)}\right)$$

## Exercises

1. Show that if  $G$  is connected and  $\mu_1 = \Delta(G)$ , then  $G$  is  $\Delta(G)$ -regular.
2. Prove Fact 4.