

The material in this handout is taken from: Reinhard Diestel, *Graph Theory (5th edition)*, Springer, 2017.

Given a set S and any $k \in \{2, \dots, |S|\}$, $[S]^k$ is the collection of all k -element subsets of S . So, $[\{1, 2, 3\}]^2 = \{\{1, 2\}, \{1, 3\}, \{2, 3\}\}$.

A graph $G = (V, E)$ has a finite **vertex set** V and a finite **edge set** $E \subseteq [V]^2$. We use i, j, u, v, w, x to denote vertices in V . The number $|V|$ of vertices is the **order** of G . A graph of order zero is empty, while graphs of order at most 1 are called trivial. An element of E is denoted by e or (i, j) . If $(i, j) \in E$, then i, j denote the endpoints of the edge (the order does not matter). A vertex i is **incident** with an edge e if $e = (i, j)$ for some $j \in V \setminus \{i\}$. Two vertices i, j are **adjacent** if $(i, j) \in E$. If $E \equiv [V]^2$, then G is the complete graph (or **clique**) on n vertices, denoted by K_n . Note that G has no self-loops (i, i) because $(i, i) \notin [V]^2$. Moreover, there can be at most one edge in G between any two pair of vertices. Such graphs are often called **simple**.

If $G' = (V', E')$ is such that $V' \subseteq V$ and $E' \subseteq [V']^2 \cap E$, then G' is a **subgraph** of G . If a subgraph G' is such that $E' \equiv [V']^2 \cap E$, then G' is called the subgraph induced by V' and denoted by $G[V']$.

The complement of a graph $G = (V, E)$ is the graph $\overline{G} = (V, \overline{E})$ such that $e \in E$ if and only if $e \notin \overline{E}$ for all $e \in [V]^2$.

Degrees. The **neighborhood** of a vertex v of G is the set $N(v)$ of vertices that are adjacent to v . The **degree** $d(v)$ of v is the cardinality of $N(v)$. A vertex with degree zero is **isolated**. The numbers $\delta(G) = \min\{d(v) : v \in V\}$ and $\Delta(G) = \max\{d(v) : v \in V\}$ are the minimum and maximum degree of G . If $\delta(G) = \Delta(G) = k$ then G is **k -regular**.

The **average degree** of G is

$$d(G) = \frac{1}{|V|} \sum_{v \in V} d(v)$$

and we obviously have $\delta(G) \leq d(G) \leq \Delta(G)$. A related quantity is the **edge density** $\varepsilon(G) = |E|/|V|$. Note that

$$|E| = \frac{1}{2} \sum_{v \in V} d(v) = \frac{1}{2} d(G) |V|$$

implying $\varepsilon(G) = d(G)/2$.

Fact 1 *The number of vertices of odd degree is always even in any graph.*

PROOF. Since $|E|$ is integer and $|E| = \frac{1}{2} \sum_{v \in V} d(v)$, then $\sum_{v \in V} d(v)$ must be even. Therefore, the number of vertices with odd degree must be even. \square

We already know that the edge density is half the average degree. Now note that the minimum degree can be larger than the edge density. For instance, in K_2 we have $\delta(G) = 1$ and $\varepsilon(G) = \frac{1}{2}$.

Fact 2 Every G with at least one edge has an induced subgraph H such that $\delta(H) > \varepsilon(H) \geq \varepsilon(G)$.

PROOF. Construct a sequence of nested subgraphs $G \equiv G_0, G_1, \dots$ induced by the vertex sets $V = V_0 \supseteq V_1 \supseteq V_2 \dots$ as follows. If V_i has a vertex v_i of degree $d(v_i) \leq \varepsilon(G_i)$ then $V_{i+1} \equiv V_i \setminus \{v_i\}$. Otherwise, stop and set $H = G_i$. If G_{i+1} is created, then

$$\varepsilon(G_{i+1}) = \frac{|E_{i+1}|}{|V_{i+1}|} = \frac{|E_i| - d(v_i)}{|V_i| - 1} \geq \frac{|E_i| - \varepsilon(G_i)}{|V_i| - 1} = \frac{|E_i|}{|V_i|} = \varepsilon(G_i)$$

The procedure stops before emptying the graph because $\varepsilon(K_1) = 0 < \varepsilon(G)$. When the procedure stops (say at $H \equiv G_k$ for some $k \geq 0$), it must be that $\delta(H) > \varepsilon(H) \geq \varepsilon(G)$, concluding the proof. \square

Paths and cycles. A **path** in $G = (V, E)$ of length $k \geq 0$ is a subgraph P_k containing $k + 1$ distinct vertices $v_0, \dots, v_k \in V$ and k edges $e_1, \dots, e_k \in E$ such that $e_i = (v_{i-1}, v_i)$ for $i = 1, \dots, k$. If $k = 0$ then $P_0 = K_1$. A **cycle** C_k in G of length $k \geq 3$ is formed when a path P_{k-1} can be extended in G to include the edge $(v_{k-1}, v_0) \in E$. The length of a shortest cycle in G is the **girth** $g(G)$, while the length of a longest cycle in G is the **circumference**. A **chord** is any edge between two vertices of a cycle which is not itself an edge of the cycle.

If a graph has a large minimum degree, then it contains long paths and cycles.

Fact 3 Every graph G with $\delta(G) \geq 2$ contains a path of length $\delta(G)$ and a cycle of length at least $\delta(G) + 1$.

PROOF. Let v_0, \dots, v_k be the vertices on any longest path P_k in G . Then $N(v_k)$ all belong to P_k (otherwise P_k is not the longest path). Therefore, $k \geq d(v_k) \geq \delta(G)$. Now let v_i the vertex of P_k with smallest index i such that $(v_i, v_k) \in E$. Then the vertices v_i, \dots, v_k form a cycle of length at least $\delta(G) + 1$ because the degree of v_k is at least $\delta(G)$. \square

The **distance** $d(i, j)$ between two vertices i, j is the length of the shortest path between them (if no path exists between the two vertices, then their distance is infinite). The **diameter** $\text{diam}(G)$ of G is the largest distance between any two vertices in G (note that the diameter can be infinite, for example when the graph has an isolated vertex). The **radius** $\text{rad}(G)$ of a graph G is the smallest distance d such that there exists a vertex whose distance from any other vertex in G is at most d . Formally,

$$\text{rad}(G) = \min_{i \in V} \max_{j \in V} d(i, j)$$

Clearly, $\text{rad}(G) \leq \text{diam}(G)$. Also, let $x \in V$ such that $d(x, v) \leq \text{rad}(G)$ for all $v \in V$. Pick any two vertices $u, v \in V$ then $d(u, v) \leq d(u, x) + d(x, v) \leq 2 \text{rad}(G)$. This shows that $\text{diam}(G) \leq 2 \text{rad}(G)$.

Girth and diameter are related as follows.

Fact 4 Every graph G containing at least a cycle satisfies $g(G) \leq 2 \text{diam}(G) + 1$.

PROOF. Let C be the shortest cycle in G . If $g(G) \geq 2 \text{diam}(G) + 2$, then C contains at least $2 \text{diam}(G) + 2$ edges. Take any two vertices x, y at opposite extremes of C . Then x, y are connected by two paths in C , say P_1 and P_2 , whose each length is at least $\text{diam}(G) + 1$. On the other hand,

the distance between x and y in G can be at most $\text{diam}(G)$ by definition of diameter. Let P be a path joining x to y in G . Note that not all the edges of P can be in C (otherwise, C is not the shortest cycle in G). Then P together with the shortest between P_1 and P_2 forms a cycle shorter than C , and we have a contradiction. \square

Connectivity. A non-empty graph G is **connected** if any two of its vertices are linked by a path in G . A maximal connected subgraph of G is a **component** of G . Any non-empty graph corresponds to a set containing at least one component. G is **k -connected** if $|V| > k$ and for all $X \subset V$ with $|X| < k$, the subgraph induced by $V \setminus X$ is connected. In words, a graph is k -connected when we cannot disconnect the graph by removing any subset of $k - 1$ vertices. Every non-empty graph is 0-connected, and the 1-connected graphs are precisely the non-trivial connected graphs (K_1 is not 1-connected because it does not have order 2).

The largest integer k such that G is k -connected is the **connectivity** $\kappa(G)$ of G . Thus $\kappa(G) = 0$ if and only if G is disconnected or $G \equiv K_1$, and $\kappa(K_n) = n - 1$ for all $n \geq 1$.

We now relate connectivity to minimum degree and to the existence of a set of edges whose removal disconnects the graph.

Theorem 5 *If G is non-trivial, then $\kappa(G) \leq |F| \leq \delta(G)$ where F is any smallest set of edges whose removal causes the graph to disconnect.*

PROOF. Let $G = (V, E)$ be non-trivial and let v be any vertex with minimum degree $\delta(G)$. Then $|F| \leq \delta(G)$ because v can be disconnected by removing the edges that are incident with $N(v)$. We now show that $\kappa(G) \leq |F|$ by a case analysis. Let $G' = (V, E \setminus F)$.

Case 1. G has a vertex v that is not incident with an edge in F . Let C be the component of G' that contains v and consider the set V_C of vertices of C that are incident with an edge of F . If we remove these vertices, then v is disconnected from the other component of G . Hence $\kappa(G) \leq |V_C|$. On the other hand, no edge in F can have both ends in C (otherwise, F is not minimal). Therefore, $|V_C| \leq |F|$.

Case 2. All vertices of G are incident with some edge in F . Pick an arbitrary vertex v and let C be the component of G' that contains v . Some $u \in N(v)$ are such that $(v, u) \in F$. The other nodes in the neighborhood of v must belong to C and are incident with distinct edges of F (otherwise, F is not minimal). Therefore $d(v) \leq |F|$, which implies $d(v) = |F| = \delta(G)$, because we already know that $|F| \leq \delta(G)$. As removing $N(v)$ disconnects v , we conclude $\kappa(G) \leq \delta(G) = |F|$. \square

Bipartite graphs. Let $r \geq 2$ be an integer. A graph $G = (V, E)$ is called r -partite if V admits a partition into r elements such that every edge has its ends in different elements: vertices in the same partition elements must not be adjacent. Instead of 2-partite one usually says bipartite. An r -partite graph in which every two vertices from different partition elements are adjacent is called complete. Note that a bipartite graph is not necessarily connected.

Clearly, a bipartite graph cannot contain an odd cycle, a cycle of odd length. One can prove that also the converse is true.

Euler tours. A walk (resp., a closed walk) is a path (resp., cycle) whose vertices may not be all

distinct. A closed walk in a graph is an Euler tour if it traverses every edge of the graph exactly once. A graph is **Eulerian** if it admits an Euler tour.

Theorem 6 (Euler, 1736) *A connected graph is Eulerian if and only if every vertex has even degree.*

PROOF. The degree condition is clearly necessary: a vertex appearing k times in an Euler tour (or $k + 1$ times, if it is the starting and finishing vertex and as such counted twice) must have degree $2k$. Conversely, we show by induction on $|E|$ that every connected graph $G = (V, E)$ with all degrees even has an Euler tour. The induction starts trivially with $|E| = 0$. Now let $|E| \geq 1$. Since all degrees are even, we can find in G a non-trivial closed walk that contains no edge more than once (see Exercise 3). Let W be such a walk of maximal length and let F be the set of its edges. We now show that $F \equiv E$, implying that W is an Euler tour. For the purpose of contradiction, suppose that $E' = E \setminus F$ has at least an edge. For every vertex $v \in V$, an even number of the edges $\{(v, u) : u \in N(v)\}$ lies in F (because W is a closed walk containing each edge only once), so the degrees of the subgraph $G' = (V, E')$ are again all even. Since G is connected, G' has an edge e incident with a vertex on W . By the induction hypothesis, the component C of G' containing e has an Euler tour. Concatenating this with W , we obtain a closed walk in G that contradicts the maximal length of W . \square

Euler tours can be found in time $\mathcal{O}(|E|)$ using Hierholzer's algorithm.

Hamilton cycles. A Hamilton cycle is a cycle that contains all vertices. A graph is Hamiltonian if it contains a Hamilton cycle.

Unlike Euler tours, only sufficient conditions are known for the existence of Hamilton cycles.

Theorem 7 (Dirac 1952) *Every graph G with $n \geq 3$ vertices and $\delta(G) \geq n/2$ is Hamiltonian.*

PROOF. Let $G = (V, E)$ be a graph with $|V| = n \geq 3$ and $\delta(G) \geq n/2$. Then G is connected: otherwise, the degree of any vertex in the smallest component C of G would be less than $|C| \leq n/2$. Let P_k be a longest path in G containing vertices v_0, \dots, v_k . Let v_i be the left end of the edge (v_i, v_{i+1}) , and v_{i+1} its right end. By the maximality of P_k , each of the $d(v_0) \geq n/2$ neighbours of v_0 is the right end of an edge of P_k , and these $d(v_0)$ edges are distinct. Similarly, at least $n/2$ edges of P_k are such that their left end is adjacent to v_k . Since P has fewer than n edges, it has an edge (v_i, v_{i+1}) with both properties.

We claim that the cycle C on vertices $v_0, v_{i+1}, \dots, v_k, v_i, \dots, v_0$ is a Hamilton cycle of G . Indeed, since G is connected, C would otherwise have a neighbour not in C which could be used to extend P_k , violating the maximality of P_k . \square

The problem of determining whether an Hamiltonian path exists in a graph is NP-complete.

Acknowledgements. Thanks to Ivan Masnari and Andrea Rovati for flagging mistakes and typos in earlier versions of this handout.

Exercises.

1. Show that every 2-connected graph contains a cycle.

2. Show that every connected graph $G = (V, E)$ contains a path of length at least

$$\min \{2\delta(G), |V| - 1\}$$

3. Show that in every connected graph whose each vertex has even degree there exists a non-trivial closed walk that contains no edge more than once.